

RESEARCH

Open Access

# IL-13 signaling via IL-13R $\alpha_2$ triggers TGF- $\beta_1$ -dependent allograft fibrosis

Stefan M Brunner<sup>1\*</sup>, Gabriela Schiechl<sup>1</sup>, Rebecca Kesselring<sup>1</sup>, Maria Martin<sup>1</sup>, Saidou Balam<sup>1</sup>, Hans J Schlitt<sup>1</sup>, Edward K Geissler<sup>1</sup> and Stefan Fichtner-Feigl<sup>1,2</sup>

## Abstract

**Background:** Allograft fibrosis still remains a critical problem in transplantation, including heart transplantation. The IL-13/TGF- $\beta_1$  interaction has previously been identified as a key pathway orchestrating fibrosis in different inflammatory immune disorders. Here we investigate if this pathway is also responsible for allograft fibrosis and if interference with the IL-13/TGF- $\beta_1$  interaction prevents allograft fibrosis.

**Methods:** FVB or control DBA/1 donor hearts were transplanted heterotopically into DBA/1 recipient mice and hearts were explanted at day 60 and 100 post-transplantation. Cardiac tissue was examined by Masson's trichrome staining and immunohistochemistry for CD4, CD8, CD11b, IL-13, Fas ligand, matrix metalloproteinase (MMP)-1, MMP-13,  $\beta_2$ -microglobulin, and Gremlin-1. Graft-infiltrating cells were isolated and analyzed by flow cytometry. IL-13 and TGF- $\beta_1$  levels were determined by enzyme-linked immunosorbent assay (ELISA) and the amount of collagen was quantified using a Sircol assay; IL-13R $\alpha_2$  expression was detected by Western blotting. In some experiments IL-13/TGF- $\beta_1$  signaling was blocked with specific IL-13R $\alpha_2$  siRNA. Additionally, a PCR array of RNA isolated from the allografts was performed to analyze expression of multiple genes involved in fibrosis.

**Results:** Both groups survived long-term (>100 days). The allogeneic grafts were infiltrated by significantly increased numbers of CD4<sup>+</sup> ( $P < 0.0001$ ), CD8<sup>+</sup> ( $P < 0.0001$ ), and CD11b<sup>+</sup> cells ( $P = 0.0065$ ) by day 100. Furthermore, elevated IL-13 levels ( $P = 0.0003$ ) and numbers of infiltrating IL-13<sup>+</sup> cells ( $P = 0.0037$ ), together with an expression of IL-13R $\alpha_2$ , were detected only within allografts. The expression of IL-13 and IL-13R $\alpha_2$  resulted in significantly increased TGF- $\beta_1$  levels ( $P < 0.0001$ ), higher numbers of CD11b<sup>high</sup>Gr1<sup>intermediate</sup>TGF- $\beta_1$ <sup>+</sup> cells, and elevated cardiac collagen deposition ( $P = 0.0094$ ). The allograft fibrosis found in these experiments was accompanied by upregulation of multiple profibrotic genes, which was confirmed by immunohistochemical stainings of allograft tissue. Blockage of the IL-13/TGF- $\beta_1$  interaction by IL-13R $\alpha_2$  siRNA led to lower numbers of CD11b<sup>high</sup>Gr1<sup>intermediate</sup>TGF- $\beta_1$ <sup>+</sup>, CD4<sup>+</sup>, CD8<sup>+</sup>, and CD11b<sup>+</sup> cells, and prevented collagen deposition ( $P = 0.0018$ ) within these allografts.

**Conclusions:** IL-13 signaling via IL-13R $\alpha_2$  induces TGF- $\beta_1$  and causes allograft fibrosis in a murine model of chronic transplant rejection. Blockage of this IL-13/TGF- $\beta_1$  interaction by IL-13R $\alpha_2$  siRNA prevents cardiac allograft fibrosis. Thus, IL-13R $\alpha_2$  may be exploitable as a future target to reduce allograft fibrosis in organ transplantation.

**Keywords:** IL-13, IL-13R $\alpha_2$ , TGF- $\beta_1$ , Allograft fibrosis, Heart transplantation

\* Correspondence: stefan.brunner@ukr.de

<sup>1</sup>Department of Surgery, University Medical Center Regensburg, Franz-Josef-Strauss-Allee 11, Regensburg 93053, Germany  
Full list of author information is available at the end of the article

## Background

Heart transplantation is an effective therapy for chronic heart failure [1]. Recent immunosuppressive strategies have reduced acute rejection episodes and improved early cardiac graft survival [2]. However, these improvements did not ameliorate chronic allograft rejection, which remains an obstacle for better long-term heart transplant survival [3]. Chronic rejection of an allograft causes an intimal fibrosis in the vessels that leads to cardiac allograft vasculopathy [4]. Another consequence of chronic rejection and inflammation is cardiac fibrosis accompanied by increased stiffness of the heart and diminished contractility [5]. Ultimately, these fibrotic reactions can result in myocardial infarction or sudden death [4,6].

On a molecular level, fibrosis is associated with a disruption of the extracellular matrix and with deposition of extracellular collagen produced by myofibroblasts [5]. In various studies TGF- $\beta_1$  has been identified as the key cytokine orchestrating fibrosis development [7,8]. TGF- $\beta_1$  is produced by macrophages after stimulation by IL-13 via the IL-13R $\alpha_2$  in the presence of IL-4 or TNF- $\alpha$  [9]. Further studies have shown that this pathway is a key initiation point for a complex fibrotic program in chronic TNBS colitis [10]. Additionally, it has been demonstrated that TGF- $\beta_1$  inhibition ameliorates lung fibrosis, chronic allograft nephropathy, and also cardiac allograft fibrosis [11-13]. However, no effective therapy to prevent heart allograft fibrosis has been identified so far, possibly because ideal murine transplant models have been lacking to study potential targets and therapies. A mouse model for the examination of cardiac allograft fibrosis should enable long-term survival of a transplanted allograft and develop cardiac fibrosis in the setting of chronic rejection. Tanaka *et al.* have developed a transplantation model in which FVB (H-2q) donor hearts were placed into DBA/1 recipients that display a similar major histocompatibility complex (MHC) (H-2q), but different non-MHC genes (CD5, CD8a, NK1.1, and Thy-1) [14]. In this model, the heart allografts survive for up to more than 100 days without immunosuppression and developed graft coronary artery disease as the result of chronic rejection.

The present study was performed under the hypothesis that the FVB to DBA/1 model is appropriate to examine cardiac graft fibrosis. Further, we hypothesized that TGF- $\beta_1$  stimulated by IL-13 signaling through IL-13R $\alpha_2$  is responsible for this allograft fibrosis and that blockage of the pathway by IL-13R $\alpha_2$ -specific siRNA can ameliorate allograft fibrosis.

## Materials and methods

### Mice and heterotopic heart transplantation

Female DBA/1 (H-2q), FVB (H-2q), and as controls BALB/c and C57BL/6 mice, 10 to 12 weeks old, were purchased from The Jackson Laboratory (Bar Harbor,

ME, USA) and housed at our local animal care facility. Animal use adhered to institutional guidelines.

Vascularized cardiac allografts were transplanted into the abdomen using a microsurgical technique as previously described by Corry *et al.* [15]. Donor hearts were perfused via the abdominal vena cava and additionally via the aortic arch with cold 0.9% saline (3 mL each) containing 500 IE heparin. Graft function was assessed by palpation of the abdomen and rejection was defined as cessation of cardiac contractility. All donor hearts had palpable contractions at the time of recovery (60 or 100 days; acute rejection 8 days).

### IL-13R $\alpha_2$ -specific siRNA

IL-13R $\alpha_2$ -specific siRNA and control (scrambled) siRNA for use in gene silencing studies were obtained from Dharmacon (Chicago, IL, USA). The siRNA (100  $\mu$ g) was encapsulated in HVJ-E and prepared as previously described before administration by intraperitoneal injection (100  $\mu$ L) every other day [10,16]. The sequence used for the siRNA is 5'-GGAATCTAATTTACAAGGA-3'.

### Histology and immunohistochemistry

Formalin-fixed and paraffin-embedded samples were prepared and sectioned (2 to 3  $\mu$ m). Tissue sections were stained with Masson's trichrome. Frozen sections (2 to 3  $\mu$ m) were blocked with 1% BSA (Biomol, Hamburg, Germany), 10% goat serum (Sigma-Aldrich, St Louis, MO, USA), or an antibody dilution buffer. As primary antibodies, rat monoclonal anti-mouse CD11b (557395; BD, Heidelberg, Germany), CD4 (550280; BD), and CD8 antibodies (Ab25478; Abcam, Cambridge, UK), a goat polyclonal anti-mouse IL-13 antibody (AF-413-NA; R&D Systems, Minneapolis, MN, USA) and a rabbit polyclonal anti-mouse Fas ligand (Ab15285; Abcam), a rabbit polyclonal anti-mouse matrix metalloproteinase (MMP)-1 (orb101432; Biorbyt, Cambridge, UK), a rabbit polyclonal anti-mouse MMP-13 (Ab39012; Abcam), a rabbit polyclonal anti-mouse  $\beta$ 2-microglobulin (Ab87483; Abcam), and a rabbit polyclonal anti-mouse Gremlin-1 antibody (Ab90670; Abcam) were used. After staining with goat anti-rat-Fab2 (sc-3822; Santa Cruz Biotechnology, Heidelberg, Germany), donkey anti-goat-Fab2 (sc-2042; Santa Cruz), or goat anti-rabbit-Fab2 (Ab64256; Abcam) secondary antibody, sections were incubated with SensiTek HRP (ScyTec Laboratories, Logan, UT, USA) and positive signals were visualized using a 3,3'-diaminobenzidine-tetrahydrochlorhydrate (DAB) kit (Merck, Darmstadt, Germany) or AEC+ High Sensitivity Substrate Chromogen kit (Dako, Hamburg, Germany). Images were captured using an Axio Observer Z1 microscope (Carl Zeiss, Oberkochen, Germany). For quantifying graft-infiltrating leukocytes, three high power fields (HPFs; 20x magnification) were counted per slide by two independent examiners.

### Western blot analyses

Cells were lysed with radioimmunoprecipitation assay buffer and the whole cell lysates obtained were subjected to SDS-PAGE. The separated proteins obtained were transferred to a nitrocellulose membrane and immunoblotted. IL-13R $\alpha_2$  was detected by incubation with a monoclonal rat anti-mouse IL-13R $\alpha_2$  (R&D Systems), followed by incubation with horseradish peroxidase-conjugated anti-rat IgG (Invitrogen, Carlsbad, CA, USA). Membranes were developed with SuperSignal West Pico Chemiluminescent Substrate (Pierce Chemical, Dallas, TX, USA) and exposed to X-ray film.

### Collagen assay

Heart allografts were harvested on day 60 and day 100 after transplantation, and homogenized in 0.5 mol/L acetic acid containing pepsin (at a concentration of 10 mg tissue/10 mL of acetic acid solution). The resulting mixture was then incubated and stirred for 24 hours at 4°C. Total soluble collagen content of the mixture was then determined with a Sircol Collagen Assay kit (Biocolor, Carrickfergus, UK), as described by the manufacturer. Acid soluble type I collagen supplied with the kit was used to generate a standard curve.

### Cell isolation from cardiac grafts and spleens

Cardiac tissue was minced in 10 mL of RPMI 1640 medium with 10% FCS, 600 U/mL collagenase II (Roche Diagnostics, Mannheim, Germany), and deoxyribonuclease I (DNase; Sigma-Aldrich). This mixture was shaken at room temperature for 2 hours and supernatant was flushed through a 100  $\mu$ m nylon cell strainer (Schubert & Weiss, Munich, Germany). Remaining tissue was again digested in 5 mL of RPMI-collagenase-DNase solution at 37°C and strained through a 100  $\mu$ m nylon strainer. Splenic tissue was minced and strained through a 100  $\mu$ m nylon strainer. Digested cell suspensions were centrifuged for 5 minutes at 1,500 rpm (4°C). To remove red blood cells, the pellet was treated with ACK lysis buffer (Lonza Walkersville, Walkersville, MD, USA) and incubated for 2 minutes at room temperature. After centrifugation, cells were suspended in HBSS medium (Gibco, Grand Island, NY, USA) and counted.

### Flow cytometry

Cell isolates were blocked with 1% mouse serum (Dako, Glostrup, Denmark) and stained with appropriate non-overlapping conjugated monoclonal antibodies (anti-Gr1 antibody from Miltenyi Biotec, Bergisch Gladbach, Germany; all other antibodies from eBioscience, San Diego, CA, USA). Intracellular staining was carried out by first fixing and permeabilizing cells with Cytfix/Cytoperm solution (BD Pharmingen, San Diego, CA, USA). Analyses were performed using a FACSCanto II

flow cytometer (BD Biosciences, San Jose, CA, USA). Data were obtained using BD CellQuest Pro acquisition software (BD Biosciences) and analyzed via FlowJo software (Tree Star Inc, Ashland, OR, USA).

### ELISA

Heart allografts were harvested at day 60 and day 100, and graft-infiltrating cells were isolated. Isolated graft-infiltrating cells were cultured at 37°C. For IL-13, we cultured  $1 \times 10^6$  cells per 1 mL medium for 48 hours; for TGF- $\beta_1$  measurements, we cultured  $1 \times 10^5$  cells per 100  $\mu$ L medium for 24 hours. During the culture period cells were stimulated with plate-bound anti-CD3 antibody (10  $\mu$ g/mL) and soluble anti-CD28 antibody (1  $\mu$ g/mL; BD Biosciences Pharmingen) for measurement of IL-13 (R&D Systems). For determination of TGF- $\beta_1$  levels (Invitrogen) cells were stimulated with plate-bound anti-CD3 antibody (10  $\mu$ g/mL), soluble anti-CD28 antibody (1  $\mu$ g/mL), and recombinant murine IL-13 (20 ng/mL; R&D Systems). Cytokine concentrations were determined in duplicate by enzyme-linked immunosorbent assay (ELISA) according to the manufacturer's instructions. TGF- $\beta_1$  was measured in medium containing TGF- $\beta_1$ -depleted human serum.

### RNA isolation and PCR array

In heart allografts recovered on day 100 after transplantation, RNA was extracted using the RNeasy Mini Kit (Qiagen, Hilden, Germany), as described by the manufacturer. One microgram of total RNA was reverse transcribed using the AffinityScript QPCR cDNA Synthesis Kit (Agilent Technologies, Böblingen, Germany). Expression of genes relevant for fibrosis was determined with a Mouse Fibrosis RT<sup>2</sup> Profiler PCR Array (SA Biosciences, Hilden, Germany) using the LightCycler 480 Real-Time PCR System (Roche).


### Statistics

All data, unless otherwise specified, are shown as the mean  $\pm$  standard error of the mean (SEM), and were compared using a two-tailed Student's test. The level of significance was set at a probability of  $P < 0.05$ .

### Results


#### FVB allografts transplanted in DBA/1 recipients showed significantly increased infiltration by CD4<sup>+</sup>, CD8<sup>+</sup>, and CD11b<sup>+</sup> cells

To determine the number of graft-infiltrating cells, heart allografts were harvested on day 60 and day 100 after transplantation, and were stained for CD4, CD8, and CD11b. In syngeneic grafts (DBA/1 into DBA/1), low numbers of CD4<sup>+</sup> (day 60,  $16 \pm 3$  and day 100,  $9 \pm 1$  cells/HPF), CD8<sup>+</sup> (day 60,  $15 \pm 2$  and day 100,  $12 \pm 3$  cells/HPF), and CD11b<sup>+</sup> cells (day 60,  $6 \pm 1$  and day 100,  $22 \pm 5$  cells/HPF) were detected (Figure 1A,B,C,D,E,F). Allogeneic heart grafts



(FVB into DBA/1) at day 60 after transplantation showed significantly higher cell numbers of CD4<sup>+</sup> ( $63 \pm 11$  cells/HPF;  $P = 0.0007$ ), CD8<sup>+</sup> ( $121 \pm 11$  cells/HPF;  $P < 0.0001$ ), and CD11b<sup>+</sup> cells ( $31 \pm 7$  cells/HPF;  $P = 0.0045$ ) compared to grafts in the syngeneic group. The numbers of CD4<sup>+</sup> and CD11b<sup>+</sup> cells in the FVB into DBA/1 group increased further by day 100 after transplantation (day 60 versus day 100, CD4<sup>+</sup>  $P = 0.0009$ ; CD11b<sup>+</sup>  $P = 0.0124$ ), whereas the increase in the number of CD8<sup>+</sup> cells did not reach statistical significance ( $P = 0.1921$ ). In comparison to control animals at day 100 after transplantation, the allogeneic group showed significantly higher levels of CD4<sup>+</sup> ( $199 \pm 31$  cells/HPF;  $P < 0.0001$ ), CD8<sup>+</sup> ( $149 \pm 17$  cells/HPF;  $P < 0.0001$ ), and CD11b<sup>+</sup> cells ( $128 \pm 34$  cells/HPF;  $P = 0.0065$ ).

**FVB allografts transplanted into DBA/1 recipients showed significantly increased levels of IL-13, IL-13R $\alpha_2$ , and TGF- $\beta_1$**   
 To examine if TGF- $\beta_1$  stimulated by IL-13 signaling is elevated in mice receiving allogeneic transplants, IL-13 levels were measured by ELISA in supernatants of cultured allograft-infiltrating cells. Syngeneic DBA/1 heart grafts showed similar IL-13 concentrations at day 60 ( $108 \pm 13$  pg/mL) and at day 100 ( $112 \pm 12$  pg/mL) ( $P = 0.8415$ ). FVB allografts placed in DBA/1 recipients showed significantly elevated IL-13 levels at day 60 ( $187 \pm 10$  pg/mL;  $P = 0.0031$ ) and at day 100 after transplantation ( $303 \pm 23$  pg/mL;  $P = 0.0003$ ) in comparison to allogeneic grafts at the same respective time points (Figure 2A). Additionally, immunohistochemical staining for IL-13 in



FVB allografts transplanted into DBA/1 mice showed significantly increased numbers of IL-13<sup>+</sup> cells/HPF at day 60 ( $16 \pm 4$  versus  $6 \pm 2$  cells/HPF;  $P = 0.0228$ ) and at day 100 ( $96 \pm 26$  versus  $7 \pm 2$  cells/HPF;  $P = 0.0037$ ), relative to the syngeneic controls (day 60 versus day 100;  $P = 0.0083$ ; Figure 2B,C). Western blot analyses of lysates from allograft-infiltrating cells indicated detectable expression of IL-13R $\alpha_2$  only in the allogeneic FVB to DBA/1 mice, both at day 60 and at day 100 after heart transplantation (Figure 2D).

As the next step, the effector cytokine TGF- $\beta_1$  was measured by ELISA after culturing and stimulating cells isolated from the allografts. In DBA/1 mice grafted with FVB hearts, significantly elevated TGF- $\beta_1$  levels were detected at day 60 ( $88 \pm 5$  versus  $46 \pm 5$  pg/mL;  $P = 0.0010$ ) and at day 100 ( $133 \pm 6$  versus  $42 \pm 7$  pg/mL;  $P < 0.0001$ ) in comparison to the syngeneic controls, and also versus the FVB control heart transplants ( $40 \pm 8$  pg/mL;  $P = 0.0048$  and  $P = 0.0009$ , respectively; Figure 2E). In accordance with

these results, flow cytometry of graft-infiltrating cells extracted from allogeneic grafts at day 100 showed a higher percentage of CD11b<sup>high</sup>Gr1<sup>intermediate</sup>TGF- $\beta$ <sub>1</sub><sup>+</sup> cells (7.3%) than in the syngeneic controls (1.2%; Figure 2F). Furthermore, flow cytometry demonstrated that these CD11b<sup>high</sup> cells were likely the only source of TGF- $\beta$ <sub>1</sub> production in this transplantation model (Figure 2G).

#### FVB allografts transplanted into DBA/1 recipients showed significantly increased levels of collagen deposition


To prove that FVB hearts transplanted in DBA/1 mice develop fibrosis, Masson's trichrome staining was performed. In these stainings, a strong collagen deposition was found in the allogeneic grafts at day 60, with a further increase in collagen deposition by day 100 after heart transplantation. No such fibrotic


collagen deposition was observed in the syngeneic control mice (Figure 3A).

A Sircol assay was conducted to better quantify collagen levels in heart allografts. With this method, the amount of collagen was found to be significantly greater in FVB hearts placed into DBA/1 recipients at day 60 ( $105 \pm 13$  versus  $61 \pm 9$   $\mu$ g/mg allograft protein;  $P = 0.0342$ ) and at day 100 after transplantation ( $139 \pm 11$  versus  $54 \pm 12$   $\mu$ g/mg allograft protein;  $P = 0.0022$ ) compared to the DBA/1-to-DBA/1 mice and also to FVB control hearts ( $P = 0.0094$ ; Figure 3B).

#### FVB allografts transplanted into DBA/1 recipients showed upregulation of profibrotic and downregulation of antifibrotic genes

To demonstrate at an mRNA-level that allogeneic grafts have upregulated profibrotic genes, RNA was isolated





from the grafted tissues and a PCR array was performed. This PCR array, which profiles the expression of 84 key genes involved in tissue remodeling and fibrosis, revealed an upregulation of IL-13 and TGF- $\beta_1$  (as detected in the previous experiments), but also a strong upregulation of other cytokines relevant for fibrosis, including IL-1 $\alpha$ , IL-1 $\beta$ , and TNF- $\alpha$ . Further, chemokines such as Ccl3, Ccl12, Ccr2, and Cxcr4, and genes involved in epithelial-mesenchymal transition or cell adhesion such as Fas ligand,  $\beta_2$ -microglobulin, integrin- $\alpha_2$ , integrin- $\beta_6$ , MMP-1a, and MMP-13, were upregulated; in contrast, Gremlin-1 and Smad6 were downregulated (Figure 4A,B).


Additionally, immunohistochemical labeling was performed for selected targets. According to these PCR array results, FVB allografts placed into DBA/1 recipients showed strong positivity for Fas ligand, MMP-1, MMP-13, and  $\beta_2$ -microglobulin at day 100, whereas Gremlin-1 staining was stronger in the syngeneic group (Figure 5). In comparison to acutely rejected grafts (BALB/c to C57BL/6), and also to naive non-transplanted FVB hearts, this expression pattern was unique in fibrosis of FVB allografts transplanted into DBA/1 recipients.

#### Specific blockage of IL-13R $\alpha_2$ abrogates TGF- $\beta_1$ production and prevents allograft fibrosis

To investigate if allograft fibrosis depends on TGF- $\beta_1$  production stimulated by IL-13 secretion, IL-13 signaling in DBA/1 recipients grafted with FVB hearts was inhibited by intraperitoneal treatment with specific IL-

13R $\alpha_2$  siRNA or control siRNA. Flow cytometric analysis of graft-infiltrating cells from hearts harvested 100 days after transplantation showed a low percentage of CD11b<sup>high</sup>Gr1<sup>intermediate</sup>TGF- $\beta_1$ <sup>+</sup> cells (0.3%) in the siRNA-treated group compared to controls (4.2%; Figure 6A). Furthermore, quantification of collagen by the Sircol assay showed significantly less collagen levels in mice treated with specific IL-13R $\alpha_2$  siRNA compared to mice treated with control siRNA (122  $\pm$  23 versus 314  $\pm$  28  $\mu$ g/mg allograft protein;  $P = 0.0018$ ) (Figure 6B). This was in accordance with Masson's trichrome staining, in which only small areas of collagen deposition were observed in siRNA-treated hearts, whereas extensive amounts of collagen were detected in control allografts (Figure 6C). To test whether the differences in TGF- $\beta_1$  production caused imbalances in CD4<sup>+</sup>Foxp3<sup>+</sup> regulatory T cells (Tregs), a flow cytometric analysis of cells from allografts was performed 100 days after transplantation. However, CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs were found to be present at equal levels in siRNA- and control siRNA-injected animals (15.6% versus 15.2%, respectively; Figure 6D). Immunohistochemical staining of allografts after therapy with specific IL-13R $\alpha_2$  siRNA showed levels of CD4<sup>+</sup>, CD8<sup>+</sup>, and CD11b<sup>+</sup> cells that were much lower after treatment with control siRNA, but similar to syngeneic transplanted animals (Figure 6E versus Figure 1A,B,C).

In this study we demonstrate for the first time that allograft fibrosis is caused by IL-13 signaling through the receptor IL-13R $\alpha_2$ , which consequently leads to elevated




**Figure 5** Immunohistochemical analysis of the 'fibrotic program' active in allogeneically transplanted grafts. Representative immunohistochemical stainings show strong positivity for Fas ligand, MMP-1, MMP-13, and  $\beta_2$ -microglobulin in allogeneically (FVB into DBA/1) and for Gremlin-1 in syngeneically (DBA/1 into DBA/1) transplanted hearts explanted at day 100, in comparison to naive non-transplanted FVB hearts and acutely rejected (BALB/c into C57BL/6) allografts (20x magnification). MMP, matrix metalloproteinase.

TGF- $\beta_1$  levels resulting in increased collagen deposition in heart allografts. Additionally, we show that inhibition of this pathway by siRNA specific for IL-13R $\alpha_2$  prevents allograft fibrosis.

The findings presented here that link IL-13 signaling via IL-13R $\alpha_2$  to allograft fibrosis are based on our previous studies showing that such signaling is essential in the development of inflammation-associated fibrosis [9,10,17]. These studies showed that IL-13 induces TGF- $\beta_1$  via a two-stage process involving: 1) induction of IL-13R $\alpha_2$  expression by IL-13 (or IL-4) signaling via IL-13R $\alpha_1$ , combined with TNF- $\alpha$  signaling through its receptor; and 2) IL-13 signaling via IL-13R $\alpha_2$  to induce an AP-1 variant containing c-Jun and Fra-2 that activates the TGF- $\beta_1$  promoter [9]. The importance of this pathway for development of fibrosis has been shown extensively by our group in bleomycin-induced lung fibrosis and chronic TNBS-induced colitis [10,17]. Thus, these previous studies provided the basis to investigate the importance of IL-13/TGF- $\beta_1$  signaling in the setting of allograft fibrosis.

The study presented here shows increasing levels of IL-13 and IL-13 $^+$  cells within allografts of transplanted mice, in contrast to control mice receiving syngeneic grafts. Multiple studies have demonstrated that IL-13 is essential for the development of dermal, gastrointestinal, and pulmonary fibrosis, as well as fibro-obliterative lesions found in the bronchiolitis obliterans (BO) syndrome [9,10,17-20]. Consistent with these studies, IL-13R $\alpha_2$  was detected only in the FVB allografts transplanted into DBA/1 recipients in our experiments. This receptor has been shown to link IL-13 signaling with further fibrotic downstream effects [9,21]. In follow-up, we detected elevated levels of TGF- $\beta_1$  in the mice receiving allogeneic grafts exclusively. Results from previous studies have indicated that TGF- $\beta_1$  is the key cytokine for development of allograft fibrosis in murine models and in humans, and that depletion of TGF- $\beta_1$  can prevent allograft fibrosis [7,12,22]. Other cytokines such as IL-6 and IL-17 can modulate the TGF- $\beta_1$ -mediated fibrotic reactions [7,8]. Additionally, a study by Faust *et al.* concluded that T cell TGF- $\beta$  signaling was required



for the development of allograft fibrosis [8]. In parallel to the elevated levels of TGF- $\beta_1$ , we found increased allograft-infiltration with CD11b<sup>high</sup>Gr1<sup>intermediate</sup>TGF- $\beta_1$ <sup>+</sup> cells in the DBA/1 mice transplanted with FVB allografts; it has been shown by our group and others that CD11b<sup>high</sup>Gr1<sup>intermediate</sup> cells are the main source for TGF- $\beta_1$  production [23-25]. In the allogeneic situation of the mouse model we used, activation of the profibrotic IL-13/TGF- $\beta_1$  interaction led to allograft fibrosis that was continuously increasing over time after transplantation.

Another important finding from this study is that allograft fibrosis can be prevented by blockage of the IL-13/TGF- $\beta_1$  interaction through specific IL-13R $\alpha_2$  siRNA. After

treatment with IL-13R $\alpha_2$  siRNA, an almost complete reduction of TGF- $\beta_1$  production by CD11b<sup>high</sup>Gr1<sup>intermediate</sup> cells (the main producers of TGF- $\beta_1$  in this model) was observed [24,25]. The reduction of TGF- $\beta_1$ -producing cells and reduced TGF- $\beta_1$  levels consequently led to diminished collagen deposition in heart allografts and therefore reduced allograft fibrosis. Tregs were also considered as contributors to the TGF- $\beta_1$  effect. While CD4<sup>+</sup>Foxp3<sup>+</sup> Tregs can produce TGF- $\beta_1$  to mediate their tolerogenic functions and expand induced regulatory T cells (iTregs), there was no difference in their numbers in control versus IL-13R $\alpha_2$  siRNA-injected mice [26-29]. Notably, after therapy with IL-13R $\alpha_2$  siRNA, CD4<sup>+</sup> and CD8<sup>+</sup> cells were

found at levels that were similar to mice receiving syngeneic grafts, and were much lower than in allotransplanted mice not given IL-13R $\alpha_2$  siRNA treatment.

For our investigations, we used a heterotopic murine heart transplantation model in which FVB hearts were placed in DBA/1 recipients. This chronic rejection model with minor multiple non-MHC mismatches has been used previously to study graft coronary artery disease [14]. We show that the FVB to DBA/1 model can also be used to examine allograft fibrosis. Over time, transplanted allografts are infiltrated by increasing numbers of CD4<sup>+</sup>, CD8<sup>+</sup>, and CD11b<sup>+</sup> cells, a fact that was also observed by Tanaka *et al.* in the original description of this transplantation model [14]. We further demonstrated by PCR array that a 'fibrotic program' is active in this FVB to DBA/1 model. Profibrotic factors such as IL-1 $\alpha$  and IL-1 $\beta$  that play a role in liver fibrosis development, and TNF- $\alpha$  which is an essential cofactor of IL-13 to induce the expression of IL-13R $\alpha_2$ , were upregulated. Further, Ccl-12 and Cxcr-4 that were both shown to be involved in pulmonary fibrosis, Ccl-3 which has been described to be important in systemic sclerosis, and Ccr-2 that is associated with allograft fibrosis, were overexpressed [9,30-33]. Additionally, the PCR array showed upregulation of other influential molecules such as Fas/Fas ligand, which is important in the development of fibrotic lesions associated with adult respiratory distress syndrome (ARDS). MMP-1 and MMP-13 (involved in remodeling processes occurring during fibrosis) and  $\beta$ 2-microglobulin were also overexpressed in the PCR array and positive in the immunohistochemistry of FVB allografts transplanted into DBA/1 recipients [34-36]. In contrast, genes like Gremlin-1 that may contribute to reversibility of lung fibrosis in rats, and Smad6 which in complex with Smurf-1 effectively attenuated TGF- $\beta_1$  signaling, were downregulated [37,38]. Altogether, these findings support the fact that this FVB to DBA/1 transplantation model is suitable not only to study graft coronary artery disease, but also to examine organ allograft fibrosis.

## Conclusions

In conclusion, this study shows that IL-13 signaling via IL-13R $\alpha_2$  induces TGF- $\beta_1$  and causes allograft fibrosis in a chronic transplant rejection model that is now also established as a model to study allograft fibrosis. Further, we demonstrate in this study that blockage of this IL-13/TGF- $\beta_1$  interaction by IL-13R $\alpha_2$  siRNA prevents heart allograft fibrosis. Together, our results indicate that IL-13R $\alpha_2$  may be exploitable as a future target to reduce allograft fibrosis in organ transplantation.

## Abbreviations

ACK: Ammonium-chloride-potassium; ARDS: Adult respiratory distress syndrome; BO: Bronchiolitis obliterans; BSA: Bovine serum albumin; DAB: 3,3'-diaminobenzidine; DNase: Deoxyribonuclease I; ELISA: Enzyme-linked

immunosorbent assay; FCS: Fetal calf serum; HBSS: Hanks' balanced salt solution; HPF: High power field; HVJ-E: Hemagglutinating virus of Japan envelope; IgG: Immunoglobulin G; IL-13: Interleukin 13; iTreg: Induced regulatory T cell; MHC: Major histocompatibility complex; MMP: Matrix metalloproteinase; PCR: Polymerase chain reaction; RPMI: Roswell park memorial institute; SEM: Standard error of the mean; siRNA: Small interfering RNA; TGF- $\beta_1$ : Transforming growth factor beta 1; TNBS: 2,4,6-trinitrobenzene sulfonic acid; TNF: Tumor necrosis factor; Treg: Regulatory T cell.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

SMB designed the study concept, collected and analyzed data, and wrote the manuscript. GS, RK, SB, and MM collected and analyzed data. HJS and EKG analyzed data and reviewed the manuscript. SFF designed the study concept, collected and analyzed data, and reviewed the manuscript. All authors read and approved the final manuscript.

## Acknowledgments

This study was supported by the University of Regensburg, Regensburg, Germany.

## Author details

<sup>1</sup>Department of Surgery, University Medical Center Regensburg, Franz-Josef-Strauss-Allee 11, Regensburg 93053, Germany. <sup>2</sup>Regensburg Center of Interventional Immunology, University Medical Center Regensburg, Regensburg, Germany.

Received: 30 May 2013 Accepted: 10 October 2013

Published: 22 October 2013

## References

- Booth AJ, Csencsits-Smith K, Wood SC, Lu G, Lipson KE, Bishop DK: Connective tissue growth factor promotes fibrosis downstream of TGF $\beta$  and IL-6 in chronic cardiac allograft rejection. *Am J Transplant* 2010, **10**:220-230.
- Guethoff S, Meiser BM, Groetzner J, Eifert S, Grinninger C, Ueberfuhr P, Reichart B, Hagl C, Kaczmarek I: Ten-year results of a randomized trial comparing tacrolimus versus cyclosporine a in combination with mycophenolate mofetil after heart transplantation. *Transplantation* 2013, **95**:629-634.
- Savasta M, Lentini S: Immunology insights into cardiac allograft rejection. *Rev Cardiovasc Disord* 2011, **12**:e68-e76.
- Huibers M, De Jonge N, Van Kuik J, Koning ES, Van Wichen D, Dullens H, Schipper M, De Weger R: Intimal fibrosis in human cardiac allograft vasculopathy. *Transpl Immunol* 2011, **25**:124-132.
- Pichler M, Rainer PP, Schauer S, Hoefler G: Cardiac fibrosis in human transplanted hearts is mainly driven by cells of intracardiac origin. *J Am Coll Cardiol* 2012, **59**:1008-1016.
- Suzuki J, Isobe M, Morishita R, Nagai R: Characteristics of chronic rejection in heart transplantation: important elements of pathogenesis and future treatments. *Circ J* 2010, **74**:233-239.
- Booth AJ, Bishop DK: TGF- $\beta$ , IL-6, IL-17 and CTGF direct multiple pathologies of chronic cardiac allograft rejection. *Immunotherapy* 2010, **2**:511-520.
- Faust SM, Lu G, Marini BL, Zou W, Gordon D, Iwakura Y, Laouar Y, Bishop DK: Role of T cell TGF $\beta$  signaling and IL-17 in allograft acceptance and fibrosis associated with chronic rejection. *J Immunol* 2009, **183**:7297-7306.
- Fichtner-Feigl S, Strober W, Kawakami K, Puri RK, Kitani A: IL-13 signaling through the IL-13 $\alpha$ 2 receptor is involved in induction of TGF- $\beta$ 1 production and fibrosis. *Nat Med* 2006, **12**:99-106.
- Fichtner-Feigl S, Young CA, Kitani A, Geissler EK, Schlitt HJ, Strober W: IL-13 signaling via IL-13R  $\alpha$ 2 induces major downstream fibrogenic factors mediating fibrosis in chronic TNBS colitis. *Gastroenterology* 2008, **135**:2003-2013. 2013 e1-7.
- Calabrese F, Kipar A, Lunardi F, Balestro E, Perissinotto E, Rossi E, Nannini N, Marulli G, Stewart JP, Rea F: Herpes virus infection is associated with vascular remodeling and pulmonary hypertension in idiopathic pulmonary fibrosis. *PLoS One* 2013, **8**:e55715.

12. Faust SM, Lu G, Wood SC, Bishop DK: **TGFbeta neutralization within cardiac allografts by decorin gene transfer attenuates chronic rejection.** *J Immunol* 2009, **183**:7307–7313.
13. Harris S, Coupes BM, Roberts SA, Roberts IS, Short CD, Brenchley PE: **TGF-beta1 in chronic allograft nephropathy following renal transplantation.** *J Nephrol* 2007, **20**:177–185.
14. Tanaka M, Zwierzchowiecka M, Mokhtari GK, Terry RD, Balsam LB, Robbins RC, Fedoseyeva EV: **Progression of alloresponse and tissue-specific immunity during graft coronary artery disease.** *Am J Transplant* 2005, **5**:1286–1296.
15. Corry RJ, Winn HJ, Russell PS: **Primarily vascularized allografts of hearts in mice. The role of H-2D, H-2K, and non-H-2 antigens in rejection.** *Transplantation* 1973, **16**:343–350.
16. Shimamura M, Morishita R, Endoh M, Oshima K, Aoki M, Waguri S, Uchiyama Y, Kaneda Y: **HVJ-envelope vector for gene transfer into central nervous system.** *Biochem Biophys Res Comm* 2003, **300**:464–471.
17. Fichtner-Feigl S, Fuss IJ, Young CA, Watanabe T, Geissler EK, Schlitt HJ, Kitani A, Strober W: **Induction of IL-13 triggers TGF-beta1-dependent tissue fibrosis in chronic 2,4,6-trinitrobenzene sulfonic acid colitis.** *J Immunol* 2007, **178**:5859–5870.
18. Aliprantis AO, Wang J, Fathman JW, Lemaire R, Dorfman DM, Lafaytis R, Glimcher LH: **Transcription factor T-bet regulates skin sclerosis through its function in innate immunity and via IL-13.** *Proc Natl Acad Sci USA* 2007, **104**:2827–2830.
19. Jakubzick C, Choi ES, Joshi BH, Keane MP, Kunkel SL, Puri RK, Hogaboam CM: **Therapeutic attenuation of pulmonary fibrosis via targeting of IL-4- and IL-13-responsive cells.** *J Immunol* 2003, **171**:2684–2693.
20. Keane MP, Gomperts BN, Weigt S, Xue YY, Burdick MD, Nakamura H, Zisman DA, Ardehali A, Saggari R, Lynch JP 3rd, Hogaboam C, Kunkel SL, Lukacs NW, Ross DJ, Grusby MJ, Strieter RM, Belperio JA: **IL-13 is pivotal in the fibro-obliterative process of bronchiolitis obliterans syndrome.** *J Immunol* 2007, **178**:511–519.
21. Strober W, Kitani A, Fichtner-Feigl S, Fuss IJ: **The signaling function of the IL-13Ralpha2 receptor in the development of gastrointestinal fibrosis and cancer surveillance.** *Curr Mol Med* 2009, **9**:740–750.
22. Rahmutola D, Marcus GM, Wilson EE, Ding CH, Xiao Y, Paquet AC, Barbeau R, Barczak AJ, Erle DJ, Olgin JE: **Molecular basis of selective atrial fibrosis due to overexpression of transforming growth factor-beta1.** *Cardiovasc Res* 2013, **99**:769–779.
23. Yang L, Huang J, Ren X, Gorska AE, Chytil A, Aakre M, Carbone DP, Matrisian Lynn M, Richmond A, Lin PC, Moses HL: **Abrogation of TGFβ signaling in mammary carcinomas recruits Gr-1+CD11b+ myeloid cells that promote metastasis.** *Cancer Cell* 2008, **13**:23–35.
24. Fichtner-Feigl S, Terabe M, Kitani A, Young CA, Fuss I, Geissler EK, Schlitt HJ, Berzofsky JA, Strober W: **Restoration of tumor immunosurveillance via targeting of interleukin-13 receptor-alpha 2.** *Cancer Res* 2008, **68**:3467–3475.
25. Terabe M, Matsui S, Park J-M, Mamura M, Noben-Trauth N, Donaldson DD, Chen W, Wahl SM, Ledbetter S, Pratt B, Letterio JJ, Paul WE, Berzofsky JA: **Transforming growth factor-β production and myeloid cells are an effector mechanism through which CD1d-restricted T cells block cytotoxic T lymphocyte-mediated tumor immunosurveillance: abrogation prevents tumor recurrence.** *J Exp Med* 2003, **198**:1741–1752.
26. Fu S, Zhang N, Yopp AC, Chen D, Mao M, Zhang H, Ding Y, Bromberg JS: **TGF-beta induces Foxp3+ T-regulatory cells from CD4+ CD25- precursors.** *Am J Transplant* 2004, **4**:1614–1627.
27. Lan Q, Zhou X, Fan H, Chen M, Wang J, Ryffel B, Brand D, Ramalingam R, Kiela PR, Horwitz DA, Liu Z, Zheng SG: **Polyclonal CD4+ Foxp3+ Treg cells induce TGFbeta-dependent tolerogenic dendritic cells that suppress the murine lupus-like syndrome.** *J Mol Cell Biol* 2012, **4**:409–419.
28. Tran DQ: **TGF-beta: the sword, the wand, and the shield of FOXP3(+) regulatory T cells.** *J Mol Cell Biol* 2012, **4**:29–37.
29. Zheng SG, Wang J, Horwitz DA: **Cutting edge: Foxp3+CD4+CD25+ regulatory T cells induced by IL-2 and TGF-beta are resistant to Th17 conversion by IL-6.** *J Immunol* 2008, **180**:7112–7116.
30. Bandinelli F, Del Rosso A, Gabrielli A, Giacomelli R, Bartoli F, Guiducci S, Matucci Cerinic M: **CCL2, CCL3 and CCL5 chemokines in systemic sclerosis: the correlation with SSC clinical features and the effect of prostaglandin E1 treatment.** *Clin Exp Rheumatol* 2012, **30**:S44–S49.
31. Kamari Y, Shaish A, Vax E, Shemesh S, Kandel-Kfir M, Arbel Y, Olteanu S, Barshack I, Dotan S, Voronov E, Dinarello CA, Apte RN, Harats D: **Lack of interleukin-1alpha or interleukin-1beta inhibits transformation of steatosis to steatohepatitis and liver fibrosis in hypercholesterolemic mice.** *J Hepatol* 2011, **55**:1086–1094.
32. Makino H, Aono Y, Azuma M, Kishi M, Yokota Y, Kinoshita K, Takezaki A, Kishi J, Kawano H, Ogawa H, Uehara H, Izumi K, Sone S, Nishioka Y: **Antifibrotic effects of CXCR4 antagonist in bleomycin-induced pulmonary fibrosis in mice.** *J Med Invest* 2013, **60**:127–137.
33. Nagai T, Tanaka M, Hasui K, Shirahama H, Kitajima S, Yonezawa S, Xu B, Matsuyama T: **Effect of an immunotoxin to folate receptor beta on bleomycin-induced experimental pulmonary fibrosis.** *Clin Exp Immunol* 2010, **161**:348–356.
34. Lopez AD, Avasarala S, Grewal S, Murali AK, London L: **Differential role of the Fas/Fas ligand apoptotic pathway in inflammation and lung fibrosis associated with reovirus 1/L-induced bronchiolitis obliterans organizing pneumonia and acute respiratory distress syndrome.** *J Immunol* 2009, **183**:8244–8257.
35. Sato M, Hwang DM, Guan Z, Yeung JC, Anraku M, Wagnetz D, Hirayama S, Waddell TK, Liu M, Keshavjee S: **Regression of allograft airway fibrosis: the role of MMP-dependent tissue remodeling in obliterative bronchiolitis after lung transplantation.** *Am J Pathol* 2011, **179**:1287–1300.
36. Zhao T, Zhao W, Chen Y, Li V, Meng W, Sun Y: **Platelet-derived growth factor-D promotes fibrogenesis of cardiac fibroblasts.** *Am J Physiol Heart Circ Physiol* 2013, **304**:H1719–H1726.
37. Farkas L, Farkas D, Gauldie J, Warburton D, Shi W, Kolb M: **Transient overexpression of Gremlin results in epithelial activation and reversible fibrosis in rat lungs.** *Am J Respir Cell Moll Biol* 2011, **44**:870–878.
38. Wang S, Sun A, Li L, Zhao G, Jia J, Wang K, Ge J, Zou Y: **Up-regulation of BMP-2 antagonizes TGF-beta1/ROCK-enhanced cardiac fibrotic signalling through activation of Smurf1/Smad6 complex.** *J Cell Mol Med* 2012, **16**:2301–2310.

doi:10.1186/2047-1440-2-16

Cite this article as: Brunner et al.: IL-13 signaling via IL-13R<sub>α2</sub> triggers TGF-β<sub>1</sub>-dependent allograft fibrosis. *Transplantation Research* 2013 **2**:16.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

